

NABIR

Biostimulation of Microbial Metabolism in Acidic FRC Subsurface Sediments

Joel E. Kostka

Florida State University





Outline

- Introduction
- Microcosm Experiments
 - Electron acceptor utilization
 - Electron donor metabolism
 - Pathways
- Conclusions
- Future work

Intro to FRC

- FRC = harsh environment for microorganisms;
Ex. pHs often 3-4 in contaminated areas
- Upon addition of electron donor and pH neutralization, extensive nitrate and metal reduction have been observed
- Thus, communities believed to be limited by: low C, pH and high nitrate, toxic metals
- What we don't know... a great deal.

FeRB and SRB catalyze the direct (enzymatic) and indirect (abiotic) reduction of U(VI)



SRB
FeRB

Populations capable of reducing metals, nitrate, halogenated compounds largely overlap



Abiotic reaction

Abiotic reaction



SRB

FeRB



Objectives

- Experimentally determine the controls or mechanisms of microbial metabolism under close to *in situ* conditions in microcosms
- Determine rates and pathways of dominant terminal-electron-accepting processes
- Compare electron donors for their biostimulation potential
- Provide inputs for reactive transport modeling to improve U bioremediation strategies

Biostimulation Experiments - Hypotheses

- For bioremediation (immobilization) of U(VI), environmental conditions of subsurface must be altered to stimulate U(VI) reduction
- U(VI) reduction rate is most likely limited by pH and the abundance of nitrate, electron donor
- Upon pH neutralization and electron donor addition, indigenous microbial communities will be stimulated to reduce NO_3^- / Fe(III)/ Mn(IV)/ U(VI)

Objectives of *In situ* Biostimulation Experiment

- To determine structure/ function relationships of metal-reducing bacteria and competing heterotrophs during *in situ* bioremediation in acidic subsurface environments
- Quantify microbial activity using geochemical analysis of groundwater/ sediments (push-pull activity tests)
- In parallel, quantify the change in the abundance/ diversity of sedimentary microbial communities using cultivation-independent methods
 - Quantitative MPN (most probable number)-PCR
 - Cloning and sequencing of 16S rRNA genes

In Situ Biostimulation using Push-Pull Activity Tests Jack Istok - OSU

- Biostimulation: addition of electron donors to increase microbial activity
- Push-pull activity tests: wells were injected with site groundwater, bicarbonate, an inert tracer, and an electron donor (glucose or ethanol)
- Groundwater chemistry was monitored over time to determine the kinetics of electron donor and acceptor utilization
- Sediment cores collected in the zone of influence surrounding wells, before and after electron donor addition

Treatments - Sediment Microcosms

- Area 1 sediments - pH 4, pH 5-7
- Neutralization with bicarbonate
- Washing to remove toxic metals
- Area 1 groundwater or distilled water
- Variety of electron donors

Initial Sediment Geochemistry

Sample	pH	Nitrate ($\mu\text{mol}/\text{cm}^3$)	Fe-HCl ($\mu\text{mol}/\text{cm}^3$)	Fe-Dithionite ($\mu\text{mol}/\text{cm}^3$)
30	4.2	22.7-31.5	15.2	414.4
31	4.8	0.1-91.7	10.8	376.4
32	4.4	0.02-17.2	18.9	407.9
33	3.6	13.6-295	14.5	467.7
34	3.8	3.3-70.5	16.3	325.0

Incubations at Bitök

Slurry	pH	Nitrate (mM)	EDs
1	5.5-6	0-0.1	Ctrl, Lac, Glu, EtOH, H ₂
2	5.5-6	20-25	Ctrl, Lac, Glu, EtOH, H ₂
3	4.3	20-25	Ctrl, Glu, EtOH
4	7.0	20-25	Ctrl, Glu, EtOH
5	6.8	1-2.5	Ctrl, Glu, EtOH
6	7.0	1-2.5	Ctrl, Glu, EtOH

Analytical Measurements

- Gases - CO_2 , H_2 , CH_4 , N_2
- Carbon substrates (HPLC)
- Nitrate
- Nitrite
- Fe(II)
- Sulfate
- pH

Area 1 Slurries







- Biostimulation clearly demonstrated
- Activity and geochemistry varies even within core
- Microbial groups display similar heterogeneity?



Slurry 2
(neutral sediment + nitrate)

Day 35


Sed. pH 5-6 - TEAPs




✓ Complete denitrification followed by substantial Fe(III) reduction

Sed. pH 5-6 - Donors


Slurry 2 Glucose-HPLC




Slurry 2 Ethanol- HPLC



Slurry 2 Lactate -HPLC



Slurry 2 - H2



✓ Glucose oxidized to acetate; ethanol completely oxidized; lactate and hydrogen not as effective electron donors

Slurry 3

Acidic, Acidic neut
washed, washed neut


Glucose

Acidic

Acidic Neut


washed

washed neut




Sed. pH 4 - TEAPs


Slurry 3 - Acidic




Slurry 4 - Acidic Neutralized



Slurry 3 - Acidic




Slurry 4 - Acidic Neutralized




✓ Activity at pH 7 but not at pH 4; nitrate reduction dominates; no Fe(III) reduction coupled to ethanol oxidation

Sed. pH 4 - Donors


Slurry 3 - Acidic - Glucose




Slurry 4 - Acidic, Neutralized - Glucose



Slurry 3 - Acidic - Ethanol




Slurry 4 - Acidic, Neutralized - Ethanol




✓ Activity at pH 7 but not at pH 4; acetate is main product

Nitrate Reduction at High Nitrate Levels (Sed. pH 4 or 5.5)


61-01-00 - Nitrate




61-03-00 - Nitrate



61-01-24 - Nitrate



61-04-00 - Nitrate



✓ Rates similar at 25 mM and 400 mM initial nitrate concentrations (pH 7)




Unamend + Ethanol

+ Glucose

*FRC, 3 wk incubation
Area 1, pH 6-7, 8/03*





Conclusions

- Metabolism inhibited under acidic conditions (pH 4)
- Electron donor/ acceptor appear to be secondary controls to pH
- FRC subsurface contains some as yet unidentified indigenous electron donor/ carbon substrate
- > sediment pH 5 or upon neutralization, classic microbial food web operates
- Nitrate and Fe(III) reduction are dominant TEAPs

Conclusions

- Electron donors vary in their biostimulation potential:
Glucose > Ethanol > Lactate > Hydrogen
- Pathways (along with activity, rates) also vary according to initial sediment pH and electron donor
- Acidic conditions shift nitrate-reducers toward incomplete reduction pathways, away from denitrification
- High nitrate concentrations do not inhibit metabolism in neutralized acidic sediments
- Ethanol did not stimulate substantial Fe(III) reduction in neutralized acidic sediments

Suggestions for Future Work

- Quantify U(VI) reduction in microcosm samples
- Quantify change in Fe mineralogy
- Identify “active” members of microbial communities
- Further characterization/ quantification of environmental parameters limiting metabolism (Ni, Al)

Acknowledgements

- Lainie Petrie, Sean O'Brien
- Special thanks to Harold Drake, Kirsten Küsel, and BITÖK
- Collaborators: David Balkwill, Joe Stucki
- FRC staff
- EU-US Task Force on Biotech. Transatlantic Environ. Biotech. Fellowship (Judy Wall, Joe Suflita, Anna Palmisano)
- The NABIR program, Biological and Environmental Research (BER), U.S. DOE (grant DE-FG02-DOER62986)

